

Mathematical Methods for Engineers (MA 713)  
Problem Sheet - 10

Change of Coordinate Matrix

- Label the following statements as true or false.
  - Suppose that  $\beta = \{x_1, x_2, \dots, x_n\}$  and  $\beta' = \{x'_1, x'_2, \dots, x'_n\}$  are ordered bases for a vector space and  $Q$  is the change of coordinate matrix that changes  $\beta'$ -coordinates into  $\beta$ -coordinates. Then the  $j$ th column of  $Q$  is  $[x_j]_{\beta'}$ .
  - Every change of coordinate matrix is invertible.
  - Let  $T$  be a linear operator on a finite-dimensional vector space  $V$ , let  $\beta$  and  $\beta'$  be ordered bases for  $V$ , and let  $Q$  be the change of coordinate matrix that changes  $\beta'$ -coordinates into  $\beta$ -coordinates. Then  $[T]_{\beta} = Q[T]_{\beta'}Q^{-1}$ .
  - The matrices  $A, B \in M_{n \times n}(F)$  are called similar if  $B = Q^t A Q$  for some  $Q \in M_{n \times n}(F)$ .
  - Let  $T$  be a linear operator on a finite-dimensional vector space  $V$ . Then for any ordered bases  $\beta$  and  $\gamma$  for  $V$ ,  $[T]_{\beta}$  is similar to  $[T]_{\gamma}$ .
- For each of the following pairs of ordered bases  $\beta$  and  $\beta'$  for  $\mathbb{R}^2$ , find the change of coordinate matrix that changes  $\beta'$ -coordinates into  $\beta$ -coordinates.
  - $\beta = \{e_1, e_2\}$  and  $\beta' = \{(a_1, a_2), (b_1, b_2)\}$
  - $\beta = \{(2, 5), (-1, -3)\}$  and  $\beta' = \{e_1, e_2\}$
  - $\beta = \{(-4, 3), (2, -1)\}$  and  $\beta' = \{(2, 1), (-4, 1)\}$
- For each of the following pairs of ordered bases  $\beta$  and  $\beta'$  for  $P_2(\mathbb{R})$ , find the change of coordinate matrix that changes  $\beta'$ -coordinates into  $\beta$ -coordinates.
  - $\beta = \{x^2, x, 1\}$  and  $\beta' = \{a_2x^2 + a_1x + a_0, b_2x^2 + b_1x + b_0, c_2x^2 + c_1x + c_0\}$
  - $\beta = \{x^2 - x + 1, x + 1, x^2 + 1\}$  and  $\beta' = \{x^2 + x + 4, 4x^2 - 3x + 2, 2x^2 + 3\}$
  - $\beta = \{2x^2 - x + 1, x^2 + 3x - 2, -x^2 + 2x + 1\}$  and  $\beta' = \{9x - 9, x^2 + 21x - 2, 3x^2 + 5x + 2\}$
- Let  $T$  be the linear operator on  $\mathbb{R}^2$  defined by

$$T \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} 2a + b \\ a - 3b \end{pmatrix},$$

let  $\beta$  be the standard ordered basis for  $\mathbb{R}^2$ , and let

$$\beta' = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\}.$$

Use the fact that

$$\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}^{-1} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$$

to find  $[T]_{\beta'}$ .

5. Let  $T$  be the linear operator on  $P_1(\mathbb{R})$  defined by  $T(p(x)) = p'(x)$ , the derivative of  $p(x)$ . Let  $\beta = \{1, x\}$  and  $\beta' = \{1 + x, 1 - x\}$ . Use the fact that

$$\begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}^{-1} = \begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & -\frac{1}{2} \end{pmatrix}$$

to find  $[T]_{\beta'}$ .

6. For each matrix  $A$  and ordered basis  $\beta$ , find  $[L_A]_{\beta}$ . Also, find an invertible matrix  $Q$  such that  $[L_A]_{\beta} = Q^{-1}AQ$ .

(a)  $A = \begin{pmatrix} 1 & 3 \\ 1 & 1 \end{pmatrix}$  and  $\beta = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 2 \end{pmatrix} \right\}$

(b)  $A = \begin{pmatrix} 13 & 1 & 4 \\ 1 & 13 & 4 \\ 4 & 4 & 10 \end{pmatrix}$  and  $\beta = \left\{ \begin{pmatrix} 1 \\ 1 \\ -2 \end{pmatrix}, \begin{pmatrix} 1 \\ -1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$

7. In  $\mathbb{R}^2$ , let  $L$  be the line  $y = mx$ , where  $m \neq 0$ . Find an expression for  $T(x, y)$ , where

(a)  $T$  is the reflection of  $\mathbb{R}^2$  about  $L$ .

(b)  $T$  is the projection on  $L$  along the line perpendicular to  $L$ .

8. Let  $T : V \rightarrow W$  be a linear transformation from a finite-dimensional vector space  $V$  to a finite-dimensional vector space  $W$ . Let  $\beta$  and  $\beta'$  be ordered bases for  $V$ , and let  $\gamma$  and  $\gamma'$  be ordered bases for  $W$ . Then prove that  $[T]_{\beta'}^{\gamma'} = P^{-1}[T]_{\beta}^{\gamma}Q$ , where  $Q$  is the matrix that changes  $\beta'$ -coordinates into  $\beta$ -coordinates and  $P$  is the matrix that changes  $\gamma'$ -coordinates into  $\gamma$ -coordinates.

9. Prove that if  $A$  and  $B$  are similar  $n \times n$  matrices, then  $\text{tr}(A) = \text{tr}(B)$ .

Hint: Use  $\text{tr}(AB) = \text{tr}(BA)$  and  $\text{tr}(A) = \text{tr}(A^t)$ .

10. Let  $V$  be a finite-dimensional vector space with ordered bases  $\alpha, \beta$ , and  $\gamma$ .

(a) Prove that if  $Q$  and  $R$  are the change of coordinate matrices that change  $\alpha$ -coordinates into  $\beta$ -coordinates and  $\beta$ -coordinates into  $\gamma$ -coordinates, respectively, then  $RQ$  is the change of coordinate matrix that changes  $\alpha$ -coordinates into  $\gamma$ -coordinates.

(b) Prove that if  $Q$  changes  $\alpha$ -coordinates into  $\beta$ -coordinates, then  $Q^{-1}$  changes  $\beta$ -coordinates into  $\alpha$ -coordinates.

11. Let  $V$  be a finite-dimensional vector space over a field  $F$ , and let  $\beta = \{x_1, x_2, \dots, x_n\}$  be an ordered basis for  $V$ . Let  $Q$  be an  $n \times n$  invertible matrix with entries from  $F$ . Define

$$x'_j = \sum_{i=1}^n Q_{ij}x_i \quad \text{for } 1 \leq j \leq n,$$

and set  $\beta' = \{x'_1, x'_2, \dots, x'_n\}$ . Prove that  $\beta'$  is a basis for  $V$  and hence that  $Q$  is the change of coordinate matrix changing  $\beta'$ -coordinates into  $\beta$ -coordinates.

12. Prove that if  $A$  and  $B$  are each  $m \times n$  matrices with entries from a field  $F$ , and if there exist invertible  $m \times m$  and  $n \times n$  matrices  $P$  and  $Q$ , respectively, such that  $B = P^{-1}AQ$ , then there exist an  $n$ -dimensional vector space  $V$  and an  $m$ -dimensional vector space  $W$  (both over  $F$ ), ordered bases  $\beta$  and  $\beta'$  for  $V$  and  $\gamma$  and  $\gamma'$  for  $W$ , and a linear transformation  $T : V \rightarrow W$  such that

$$A = [T]_{\beta}^{\gamma} \quad \text{and} \quad B = [T]_{\beta'}^{\gamma'}.$$

Hints: Let  $V = F^n$ ,  $W = F^m$ ,  $T = L_A$ , and  $\beta$  and  $\gamma$  be the standard ordered bases for  $F^n$  and  $F^m$ , respectively. Now apply the results of the above exercise to obtain ordered bases  $\beta'$  and  $\gamma'$  from  $\beta$  and  $\gamma$  via  $Q$  and  $P$ , respectively.